

## Research article

# Sexual inactivity and sexual satisfaction among women living with HIV in Canada in the context of growing social, legal and public health surveillance

Angela Kaida<sup>§,1</sup>, Allison Carter<sup>1,2</sup>, Alexandra de Pokomandy<sup>3,4</sup>, Sophie Patterson<sup>1,2</sup>, Karène Proulx-Boucher<sup>3</sup>, Adriana Nohpal<sup>2</sup>, Paul Sereda<sup>2</sup>, Guillaume Colley<sup>2</sup>, Nadia O'Brien<sup>3,4</sup>, Jamie Thomas-Pavanel<sup>5</sup>, Kerrigan Beaver<sup>5</sup>, Valerie J Nicholson<sup>1</sup>, Wangari Tharao<sup>6</sup>, Mylène Fernet<sup>7</sup>, Joanne Otis<sup>7</sup>, Robert S Hogg<sup>1,2</sup>, Mona Loutfy<sup>5,8</sup>, on behalf of the CHIWOS Research Team<sup>†</sup>

<sup>§</sup>**Corresponding author:** Angela Kaida, Faculty of Health Sciences, Simon Fraser University, Blusson Hall Rm 10522, 8888 University Drive, Burnaby, BC, Canada V5A 1S6. Tel: +1 778 782 9068. Fax: +1 778 782 5927. (kangela@sfu.ca)

<sup>†</sup>Additional team members are listed at the end of the manuscript.

## Abstract

**Introduction:** Women represent nearly one-quarter of the 71,300 people living with HIV in Canada. Within a context of widespread HIV-related stigma and discrimination and on-going risks to HIV disclosure, little is known about the influence of growing social, legal and public health surveillance of HIV on sexual activity and satisfaction of women living with HIV (WLWH).

**Methods:** We analyzed baseline cross-sectional survey data for WLWH ( $\geq 16$  years, self-identifying as women) enrolled in the Canadian HIV Women's Sexual and Reproductive Health Cohort Study (CHIWOS), a multisite, longitudinal, community-based research study in British Columbia (BC), Ontario (ON) and Quebec (QC). Sexual inactivity was defined as no consensual sex (oral or penetrative) in the prior six months, excluding recently postpartum women ( $\leq 6$  months). Satisfaction was assessed using an item from the Sexual Satisfaction Scale for Women. Multivariable logistic regression analysis examined independent correlates of sexual inactivity.

**Results:** Of 1213 participants (26% BC, 50% ON, 24% QC), median age was 43 years (IQR: 35, 50). 23% identified as Aboriginal, 28% as African, Caribbean and Black, 41% as White and 8% as other ethnicities. Heterosexual orientation was reported by 87% of participants and LGBTQ by 13%. In total, 82% were currently taking antiretroviral therapy (ART), and 77% reported an undetectable viral load (VL  $< 40$  copies/mL). Overall, 49% were sexually inactive and 64% reported being satisfied with their current sex lives, including 49% of sexually inactive and 79% of sexually active women ( $p < 0.001$ ). Sexually inactive women had significantly higher odds of being older (AOR = 1.06 per year increase; 95% CI = 1.05–1.08), not being in a marital or committed relationship (AOR = 4.34; 95% CI = 3.13–5.88), having an annual household income below \$20,000 CAD (AOR: 1.44; 95% CI = 1.08–1.92), and reporting high (vs. low) HIV-related stigma (AOR = 1.81; 95% CI = 1.09–3.03). No independent association was found with ART use or undetectable VL.

**Conclusions:** Approximately half of WLWH in this study reported being sexually inactive. Associations with sexual dissatisfaction and high HIV-related stigma suggest that WLWH face challenges navigating healthy and satisfying sexual lives, despite good HIV treatment outcomes. As half of sexually inactive women reported being satisfied with their sex lives, additional research is required to determine whether WLWH are deliberately choosing abstinence as a means of resisting surveillance and disclosure expectations associated with sexual activity. Findings underscore a need for interventions to de-stigmatize HIV, support safe disclosure and re-appropriate the sexual rights of WLWH.

**Keywords:** HIV; women; Canada; sexual and reproductive health; sexual abstinence; sexual satisfaction; community-based research; antiretroviral therapy; CHIWOS.

**Received** 8 May 2015; **Revised** 23 September 2015; **Accepted** 5 October 2015; **Published** 1 December 2015

**Copyright:** © 2015 Kaida A et al; licensee International AIDS Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

Globally, women account for over half of all adults living with HIV [1]. In Canada, approximately one-quarter of the 71,300 people living with HIV (PLWH) are women, nearly double the proportion observed in 1999 (12%) [2]. With early and sustained use of antiretroviral therapy (ART), women living

with HIV (WLWH) are living longer and healthier lives [3–5] with improved sexual and reproductive options accompanying lowered risks of sexual and perinatal HIV transmission [6–8]. This altered landscape of HIV risk has re-ignited global discourse regarding the need for a rights-based approach to sexual health [9–12]. Sexual health research and programming

targeting WLWH, however, are largely focused on risk behaviours (such as condom use) rather than broader sex-positive considerations of sexual intimacy, well-being and satisfaction [13,14]. Moreover, prevailing concern about individual-level behaviours ignores the broader clinical, legal and social factors that regulate the sexual lives and rights of WLWH [15,16].

The criminalization of HIV non-disclosure [17] and “Treatment as Prevention” (TasP) [18] are two legal and public health strategies aimed at preventing HIV transmission that have emerged in many global settings, including Canada [11,19–21]. While both initiatives place sexual activity of PLWH and “Positive Prevention” (HIV prevention strategies directed at PLWH) at the centre of their efforts, they are theoretically opposing and deviate from the “shared responsibility” messaging for HIV prevention endorsed by the Global Network of People Living with HIV (GNP+), the Canadian AIDS Society and other international agencies [22,23]. Growing evidence demonstrates that early and sustained use of ART for greater than six months with viral suppression generates a very low risk of HIV transmission during condomless sex between HIV sero-discordant sexual partners, with recent studies reporting a transmission risk approaching zero [24–26]. These findings have supported the implementation of TasP initiatives in a number of global jurisdictions [20,27,28], where PLWH are monitored along a care cascade from HIV diagnosis to linkage and retention in care, and initiation and adherence to ART to achieve sustained viral suppression [29,30].

Despite the evidence and growing optimism about what it means to live with HIV, in terms of improved quality of life, clinical health and lowered transmission risks, Canada has among the most aggressive judicial approaches to prevent perceived sexual exposure to HIV through the criminalization of HIV non-disclosure [19,21]. In October 2012, the Supreme Court of Canada ruled that PLWH are legally required to disclose their HIV status to sexual partners prior to sexual activity that poses a “realistic possibility” of HIV transmission [17,31]. The Supreme Court defined realistic possibility as any sexual activity without the use of a condom and without a low HIV plasma viral load (defined by the court as VL < 1500 copies/mL). PLWH who fail to meet both criteria and do not disclose their HIV status to sexual partners risk a criminal charge of aggravated sexual assault. If convicted, this charge results in jail time with a maximum sentence of life imprisonment and mandatory listing on a national Sexual Offender Registry. The inconsistency between legal definitions of the “realistic possibility” of HIV transmission and contemporary scientific assessments of HIV transmission risk and prognosis detract from rights-based approaches to improving the sexual health of WLWH and **propagate misconceptions about the sexual and reproductive realities of living with HIV.**

For many WLWH around the world, such public health and legal HIV prevention strategies have introduced increased surveillance (including monitoring VL, HIV disclosure and condom use) and present attendant consequences to their sexual lives. Importantly, expectations of HIV status disclosure exist within a context of widespread HIV-related fear, violence, and stigma and discrimination, which when combined with gender and relationship power inequities, disproportionately

compromise WLWH’s navigation of intimate relationships [32–35]. Together, these social factors mute conversations about sexuality and HIV, deny risks that WLWH face with forced disclosure and/or condom use, compromise opportunities for safe disclosure and sexual relationships for WLWH and reduce the willingness and agency of WLWH to seek out critical health services [36]. In Canada and elsewhere, increasing use of ART has not substantially alleviated the presence or impact of stigma and discrimination [37] despite attempts to normalize HIV through initiatives under the TasP umbrella [38,39].

In response to growing social, legal and public health surveillance, WLWH may adopt various strategies to protect and navigate their sexual lives, including the avoidance of sexual and romantic partnerships. Indeed, previous research has suggested that WLWH may forgo sexual activity to avoid expectations and risks of HIV disclosure to sexual partners [32–34]. To understand drivers of sexual activity decision-making among WLWH in Canada, we measured the prevalence of sexual inactivity and sexual satisfaction among a cohort of WLWH and assessed demographic, HIV clinical and socio-structural correlates of sexual inactivity.

## Methods

### Study setting

In Canada, approximately 16,600 women were living with HIV in 2011 [40], of whom 81% lived in one of the three provinces: British Columbia (BC), Ontario or Quebec.

### Study design

Baseline cross-sectional data were analyzed for WLWH enrolled in the Canadian HIV Women’s Sexual and Reproductive Health Cohort Study (CHIWOS; www.chiwos.ca), a large, multisite, longitudinal, community-based research (CBR) project conducted by, with and for WLWH in BC, Ontario and Quebec (full cohort:  $n = 1427$ ). The primary objective of CHIWOS is to assess the prevalence, barriers and facilitators to use of women-centred HIV care, and the impact of such patterns of use on health outcomes. CHIWOS is grounded in Critical Feminist theory [41] and CBR principles, and guided by a Social Determinants of Women’s Health framework [42,43] (described in detail elsewhere [44]). Consistent with CBR methodology, WLWH and allied clinicians, researchers, community partners, social and public health service delivery personnel, and policy-makers were involved in all stages of this research.

### Study population and recruitment

This analysis includes participants enrolled in CHIWOS between 27 August 2013 and 13 March 2015. WLWH aged  $\geq 16$  years from one of the three enrolling study provinces were eligible to participate. CHIWOS is inclusive to self-identified trans and cis gendered women, two-spirited women, gender queer and women of other gender identities. Given the importance of including young women’s voices in the study, an exemption was sought and approved such that parental/guardian consent for women aged 16–18 years was not required.

WLWH were recruited for participation through peer word-of-mouth, HIV clinics in large and small communities across the three provinces, AIDS Service Organizations (ASOs), non-HIV

community-based organizations (e.g. immigration and refugee services, women's shelters, harm reduction and sex worker support services), the networks of our national Steering Committee and three provincial Community Advisory Boards, and online methods (e.g. Listservs for WLWH, and our study website ([www.chiwos.ca](http://www.chiwos.ca)), Facebook page ([www.facebook.com/CHIWOS](http://www.facebook.com/CHIWOS)) and Twitter feed ([www.twitter.com/CHIWOS](http://www.twitter.com/CHIWOS) research).

Consistent with data showing an over-representation of Aboriginal and African, Caribbean and Black (ACB) women among WLWH in Canada, targeted efforts were made to recruit Aboriginal and ACB participants. These efforts included hiring a diverse national team of 38 WLWH as Peer Research Associates (PRAs), including 6 Aboriginal and 14 ACB PRAs, and the creation of a CHIWOS Aboriginal Advisory Board whose members assisted and advised on community outreach and recruitment.

Overall, 1355 women were enrolled in CHIWOS at time of analysis. For this analysis, 142 (10%) women were excluded, including 84 who requested to skip the Sexual Health section of the survey, 7 who reported a live birth at or less than six months prior to interview, 40 who reported never having had consensual sex, and 13 who were missing data on recent consensual sex (the primary outcome), yielding an analytic sample of 1213 WLWH. Given our objective to measure intentional sexual activity, women who experienced non-consensual sex in the six months prior to interview without report of consensual sexual activity, were not considered sexually active.

#### Data collection

All participants completed a structured, online questionnaire (supported by FluidSurveys™ software) at baseline (study enrolment), administered by PRAs (WLWH hired as members of the study team). PRAs underwent extensive training in research ethics, CBR, consenting, administering questionnaires, social positioning, self-care and support for participants [45]. Questionnaires were conducted in English or French and administered in-person in a confidential setting at collaborating HIV clinics, ASOs or community organizations, or in women's homes. For some participants in rural or remote areas, questionnaires were administered via phone or Skype. Median survey completion time was 89 minutes [IQR: 71, 115] and participants received a \$50 honorarium for their participation, regardless of whether all questions or sections of the questionnaire were completed.

#### Measures

A national team of experts in women's health and HIV contributed to the development of the CHIWOS questionnaire, which was designed to maximize psychometric validity and reliability [46] and collected information on socio-demographics, HIV medical history and clinical care, experiences of violence and stigma and discrimination, and sexual, reproductive, women's and emotional health outcomes. Given the sensitive nature of questions about sexual health and violence, participants were given the option of completing these survey sections with the PRA, independently, or skipping the sections entirely.

#### Primary outcomes: sexual inactivity and sexual satisfaction

The primary outcome was recent sexual inactivity (vs. sexual activity) defined as no (vs. any) consensual sex (oral or penetrative) in the six months prior to interview.

Among all women, regardless of recent sexual activity or inactivity, sexual satisfaction was assessed using a five-point Likert scale item from the Sexual Satisfaction Scale for Women (SSS-W) [47]: "Overall, how satisfactory or unsatisfactory is your present sex life." Responses were dichotomized into satisfactory ("Completely/Very/Reasonably satisfactory") vs. unsatisfactory ("Not very/Not at all satisfactory").

#### Covariates

Covariates of sexual inactivity included socio-demographic characteristics (i.e. age, education, gender identity, sexual orientation, ethnicity, annual household income, relationship status, and number of children); self-reported clinical HIV history and outcomes, including years living with HIV, receipt of HIV medical care in past year (yes vs. no), receipt of ART in past year (yes vs. no), most recent VL (undetectable (< 40 copies/mL) vs. detectable  $\geq 40$  copies/mL, shown to have a high positive predictive value [48]), most recent CD4 cell count (< 200 cells/mm<sup>3</sup> vs. 200–500 cells/mm<sup>3</sup> vs. > 500 cells/mm<sup>3</sup>), as well as having ever discussed the impact of VL on risk of HIV transmission with a healthcare provider (yes vs. no) and perceptions of how ART changes personal risk of transmitting HIV (lowers the risk of transmission vs increases the risk of transmission or makes little difference); the SF-12 physical and mental health-related quality of life summary scores (scored on a 0–100 scale, where a higher score indicates better health) [49,50]; depression (measured using the 10-item Center for Epidemiologic Studies Depression Scale (CES-D 10), scored on a 0–30 scale, with higher scores indicating greater depression symptomology and a cutoff score of  $\geq 10$  considered indicative of "probable depression") [51,52]; and HIV-related stigma measured using the 10-item HIV Stigma Scale (HSS) [53,54]. Scores for the HSS range from 0 to 100, with higher scores indicating higher stigma. Scores greater or equal to the scale median were considered "high HIV-related stigma" vs. "low HIV-related stigma."

Among sexually active women, we describe types of sexual partnerships and knowledge about HIV status. In the survey, "Regular Sexual Partners" were defined to include, but not limited to, "spouses, common law partners, long term relationships, friends with benefits, or partners seen on and off for some time." "Casual Sexual Partners" were defined to include, but not limited to, "serious sexual relationships that have recently begun, new sexual relationships that exist but you're not sure about, chance sexual encounters, or one night stands."

#### Data analysis

Descriptive statistics (median Interquartile Range [IQR]) for continuous variables and n (%) for categorical variables) were used to characterize baseline distributions of study variables. Baseline differences between sexually inactive and sexually active women were compared using Wilcoxon rank sum test for continuous variables and Pearson  $\chi^2$  or Fisher's exact test for categorical variables.

We fit a multivariable logistic regression explanatory model to the data to examine covariates of sexual inactivity. After testing normality assumptions and collinearity, variables with a significant association with sexual inactivity in bivariate analyses (at  $p < 0.20$ ) were considered for the full model to obtain the relative contribution of each covariate. Model selection was achieved by minimizing the Akaike Information Criterion while maintaining Type III  $p$ -values for covariates below 0.20 [55]. All statistical tests were two-sided and were considered statistically significant at  $\alpha = 0.05$ . Data were analyzed using SAS version 9.3 (SAS Institute, Inc., Cary, NC).

#### Ethical statement

All participants provided written, voluntary informed consent (or oral consent with a study team member present as a witness for surveys conducted by phone or Skype) at enrolment in the survey phase of the CHIWOS study. Ethical approval for all study procedures was provided by the Research Ethics Boards of Simon Fraser University, University of British Columbia/Providence Health, Women's College Hospital and McGill University Health Centre.

## Results

### Baseline characteristics

Of 1213 participants, 26% were from BC, 50% from Ontario and 24% from Quebec. Median age was 43 years [IQR: 35, 50], 95% identified as cis gender women and 5% identified as trans women, two-spirited or gender queer. Eighty-seven percent were heterosexual and 13% identified as LGBTQ (Lesbian, Gay, Bi-Sexual, Two-spirited, or Queer). Nearly one-quarter (23%) identified as Aboriginal, 28% ACB, 41% White and 8% other ethnicities. Sixty-three percent had an annual household income  $< \$20,000$  CAD, and 34% were married, common law or in a relationship.

Median years living with HIV was 10.8 [IQR: 5.9, 16.8], 94% reported receiving HIV medical care in the past year and 82% were currently taking ART. By self-report, 77% had undetectable VL and 50% had CD4 cell count  $> 500$  cells/mm<sup>3</sup> at most recent visit. Sixty-nine percent of participants reported discussing the impact of VL on HIV transmission risk with their healthcare provider, and overall, 81% perceived that use of antiretrovirals lowers HIV transmission risk. Median SF-12 physical and mental health summary scores were 48 [IQR: 33, 56] and 42 [IQR: 31, 52], respectively. Median HIV stigma score was 60 [IQR: 50, 73] with 48% classified as experiencing high (vs. low) HIV-related stigma. Median depression symptom severity score (based on CES-D) was 9 [IQR: 4, 15], with 49% meeting the criteria for probable depression (Table 1).

### Sexual inactivity, sexual satisfaction and summary of sexual partnerships

Forty-nine percent (49%) of participants reported being sexually inactive in the six months prior to baseline interview. Sixty-four percent (64%) of women reported being satisfied with their current sex lives, including 49% of sexually inactive women and 79% of sexually active women ( $p < 0.001$ ).

Of sexually active WLWH ( $n = 618$ ), 81% had one and 6% had two or more regular sexual partners in the previous six

months. Of those with at least one regular sexual partner ( $n = 543$ ), 27% reported that their primary sexual partner was HIV-positive, 64% reported that the partner was HIV-negative and for 9% the partner's HIV status was unknown. Ninety-one percent (91%) reported that their current or most recent regular partner knew the participant's HIV status at their last sexual encounter. One-fifth (21%) of all sexually active participants reported having a casual sexual partner in the previous six months.

### Correlates of sexual inactivity

In unadjusted analyses, sexual inactivity was associated (at  $p < 0.20$ ) with older age, gender identity, sexual orientation, household income  $< \$20,000$  per year, not being in a marital or committed relationship, longer time living with HIV, current ART use, poorer physical health, high HIV-related stigma scores, probable depression, sexual dissatisfaction and having discussed the role of VL on HIV transmission risk with a healthcare provider (Table 2).

In the logistic regression model, sexually inactive women had significantly higher adjusted odds of being older (AOR = 1.06 per year increase in age; 95% CI = 1.05–1.08), not being in a marital or committed relationship (AOR = 4.34; 95% CI = 3.13–5.88), having an annual household income below  $\$20,000$  CAD (AOR = 1.44; 95% CI: 1.08–1.92), and reporting high (vs. low) HIV-related stigma (AOR = 1.81; 95% CI = 1.10–3.03). No independent association was found with current ART use, undetectable plasma HIV VL or CD4  $> 500$  cells/mL. However, sexually inactive women were significantly more likely to report not having discussed the role of VL on decreasing HIV transmission risk with a healthcare provider (AOR = 1.57; 95% CI: 1.16–2.11; Table 2).

## Discussion

This is the first multisite cohort study to evaluate sexual inactivity among WLWH in Canada in the modern era of TasP and HIV criminalization. Nearly half of WLWH in this Canadian cohort reported being sexually inactive and over two-thirds report being satisfied with their sexual lives. Other studies have similarly shown that sexual inactivity is common among PLWH, particularly women. The Women's Interagency HIV Study (WIHS), an on-going cohort study of WLWH in the United States (US), found that 35% of WLWH reported no vaginal, oral or anal sex over a six-month time period compared with 23% of HIV-negative women [56]. Other US research has found that WLWH are significantly more likely to be sexually inactive (34%) compared with HIV-positive gay and bisexual men (28%) [57]. Similar results were seen in a study by Bogart *et al.* [58], where a higher proportion of WLWH (18%) were deliberately abstinent compared to gay and bisexual men with HIV (11%). The prevalence of sexual inactivity (49%) in our study was notably higher than estimates from other HIV cohorts. Some of this difference may be explained by the older median age of WLWH enrolled in CHIWOS compared with other referenced cohorts and the evidence that sexual activity and satisfaction decline with increasing age [59,60]; and shown in Table 1. Some difference may also be explained by our distinction between and exclusion of non-consensual

**Table 1. Baseline characteristics of women living with HIV enrolled in CHIWOS overall and by sexual activity (n = 1213)**

Characteristic	Overall (n = 1213)		Sexually active (n = 618)	Sexually inactive (n = 595)	p-value
	n (%) or median [IQR]	Total n	n (%) or median [IQR]		
Median age [IQR] (years)	43 [35, 50]	1213	40 [33, 46]	46 [38, 54]	<0.001
Age category (years)					
16 to <30	114 (9.4)	1213	83 (13)	31 (5.2)	<0.001
30 to <40	366 (30)		218 (35)	148 (25)	
40 to <50	392 (32)		215 (35)	177 (30)	
50+	341 (28)		102 (17)	239 (40)	
Province					
British Columbia (BC)	317 (26)	1213	172 (28)	145 (24)	0.009
Ontario	607 (50)		283 (46)	324 (55)	
Quebec	289 (24)		163 (26)	126 (21)	
Gender identity <sup>a</sup>					
cis gender woman	1157 (95)	1213	581 (94)	576 (97)	0.02
Trans woman, two-spirited, gender queer, or other gender	56 (5)		37 (6)	19 (3)	
Sexual orientation <sup>b</sup>					
Heterosexual	1050 (87)	1209	525 (85)	525 (88)	0.121
LGBTQ	159 (13)		90 (15)	69 (12)	
Ethnicity					
Aboriginal	277 (23)	1213	142 (23)	135 (23)	0.391
African, Caribbean and Black Canadian (ACB)	338 (28)		164 (27)	174 (29)	
White	502 (41)		268 (43)	234 (39)	
Other ethnicities	96 (8)		44 (7.1)	52 (8.7)	
Education					
<High school	180 (15)	1207	93 (15)	87 (15)	0.798
≥High school	1027 (85)		520 (85)	507 (85)	
Annual household income (CAD)					
≥\$20,000	419 (35)	1213	242 (39)	177 (30)	<0.001
<\$20,000	759 (63)		353 (57)	406 (68)	
Don't know/prefer not to answer	35 (3)		23 (3.7)	12 (2.0)	
Relationship status					
Single/separated/divorced/widowed	802 (66)	1207	306 (50)	496 (84)	<0.001
Married/common law/in a relationship	405 (34)		307 (50)	98 (17)	
Number of children					
0	372 (32)	1162	175 (30)	197 (34)	0.204
1–3	630 (54)		332 (57)	298 (52)	
4+	160 (14)		78 (13)	82 (14)	
Median time living with HIV [IQR] (years)	10.8 [5.9, 16.8]	1180	10.1 [5.6, 15.8]	11.7 [6.2, 17.6]	0.004
Received HIV medical care in the past year					
Yes	1139 (94)	1212	574 (93)	565 (95)	0.102
No	73 (6.0)		44 (7.1)	29 (4.9)	
Currently taking ART					
Yes	999 (82)	1213	500 (81)	499 (84)	0.176
No	214 (18)		118 (19)	96 (16)	
Current plasma viral load (self-reported)					
Detectable (≥40 copies/mL)	173 (14)	1213	91 (15)	82 (14)	0.869
Undetectable (<40 copies/mL)	939 (77)		477 (77)	462 (78)	
Don't know/never received results	101 (8.3)		50 (8.1)	51 (8.6)	



**Table 1 (Continued)**

Characteristic	Overall (n = 1213)		Sexually active (n = 618)	Sexually inactive (n = 595)	p-value
	n (%) or median [IQR]	Total n	n (%) or median [IQR]		
Current CD4 cell count (self-reported)					
<200 cells/mm <sup>3</sup>	66 (5.5)	1211	34 (5.5)	32 (5.4)	0.597
200–500 cells/mm <sup>3</sup>	329 (27)		166 (27)	163 (28)	
>500 cells/mm <sup>3</sup>	606 (50)		302 (49)	304 (51)	
Don't know/never received results	210 (17)		116 (19)	94 (16)	
Discussed VL on HIV transmission risk with healthcare provider					
Yes	826 (69)	1201	461 (75)	365 (62)	<0.001
No	375 (31)		151 (25)	224 (38)	
Perception of how ART changes HIV transmission risk					
Lowers risk of transmission	979 (81)	1213	509 (82)	470 (79)	0.214
Increases risk of transmission or makes little difference	130 (11)		57 (9.2)	73 (12)	
Don't know	104 (8.6)		52 (8.4)	52 (8.7)	
Median health-related quality of life score [IQR]					
Physical health summary score	48 [33, 56]	1206	50 [36, 56]	46 [32, 55]	0.004
Mental health summary score	42 [31, 52]	1206	42 [31, 52]	43 [32, 53]	0.537
HIV stigma scale					
High HIV-related stigma	575 (48)	1200	280 (46)	295 (50)	0.155
Low HIV-related stigma	625 (52)		330 (54)	295 (50)	
Satisfaction with present sex life					
Completely, very or reasonably satisfied	777 (64)	1213	488 (79)	289 (49)	<0.001
Not very or not at all satisfied	347 (29)		115 (19)	232 (39)	
Prefer not to answer	89 (7.3)		15 (2.4)	74 (12)	
Probable depression (CESD-10)					
No (score <10)	600 (51)	1173	330 (55)	270 (47)	0.007
Yes (score ≥10)	573 (49)		270 (45)	303 (53)	

Notes: <sup>a</sup>Gender identity was self-reported as cis or trans gender woman, two-spirited, gender queer or an 'other' gender identity. Given small numbers, we grouped participants self-identifying as trans gender, two-spirited, gender queer or 'other' gender for this analysis. Of the 56 women included in this grouped category, 48 (86%) identified as transgender women.

<sup>b</sup>LGBTQ includes participants who identify as Lesbian, Gay, Bi-sexual, Two-spirited or Queer; CAD = Canadian dollars; ART = antiretroviral therapy; VL = viral load.

sex in our definition of sexual activity. However, the remaining difference may be explained by a high reported prevalence of HIV stigma and the context of surveillance for WLWH in the Canadian setting [61]. Our estimates of sexual inactivity are also significantly higher than those reported in the general population (10–14%) [62], echoing previous research that an HIV diagnosis impacts the sexual health of women [63] even several years after the initial diagnosis [64].

The association between sexual inactivity and high HIV-related stigma and sexual dissatisfaction but not ART use or viral suppression suggests that WLWH face challenges navigating healthy, satisfying and safe sexual lives, despite good treatment outcomes. This suggests that good treatment outcomes alone do not lead to higher likelihood of sexual

activity or eliminate prevailing socio-structural barriers to sexual health for WLWH.

HIV-related stigma and discrimination are critical barriers to cultivating loving, intimate relationships, facilitating HIV status disclosure and engaging with healthcare for WLWH [65,66]. The increasingly strict use of the criminal law for HIV non-disclosure against PLWH in Canada presents an additional form of stigmatization [67] and additional challenges for WLWH initiating sexual relationships and navigating HIV disclosure to sexual partners. This judicial approach shifts the responsibility for condom use and the burden of proof onto PLWH, introducing the potential for false accusations of HIV non-disclosure by sexual partners and concerns of secondary disclosure within the wider community [68,69]. Observations

**Table 2. Unadjusted and adjusted odds ratios for correlates of sexual inactivity in the previous six months among women living with HIV enrolled in CHIWOS**

Characteristics	Sexually inactive vs. Sexually active over the last six months	
	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Age, per one year increase	1.06 (1.05–1.08)	1.06 (1.05–1.08)
Sexual orientation		
Heterosexual	1.0	
LGBTQ	0.75 (0.54–1.05)	
Annual household income (CAD)		
≥\$20,000	1.0	1.0
<\$20,000	1.57 (1.24–2.0)	1.44 (1.08–1.92)
Relationship status		
Married/common law/in a relationship	1.0	1.0
Single/separated/divorced/widowed	5.00 (3.84–6.67)	4.34 (3.13–5.88)
Currently taking ART		
Yes	1.0	
No	0.82 (0.61–1.10)	
Discussed VL on HIV transmission risk with healthcare provider		
Yes	1.0	1.0
No	1.87 (1.46–2.4)	1.57 (1.16–2.11)
HIV-related stigma		
Low HIV-related stigma	1.0	1.0
High HIV-related stigma	1.18 (0.94–1.47)	1.81 (1.09–3.03)
Probable depression (CESD-10)		
No (score < 10)	1.0	
Yes (score ≥ 10)	1.37 (1.09–1.73)	
Physical Health Summary score	0.99 (0.98–0.99)	
Satisfaction with present sex life		
Completely, very, or reasonably satisfied	1.0	
Not very or Not at all satisfied	3.41 (2.61–4.45)	
Prefer not to answer	8.33 (4.69–14.8)	

Note: Model adjusted for Province of residence; LGBTQ includes participants who identify as Lesbian, Gay, Bi-sexual, Two-spirited or Queer; CAD = Canadian dollars; ART = antiretroviral therapy; VL = viral load.

that the sexuality of PLWH is being policed [70] and reports of discriminating medical surveillance in the reproductive healthcare setting [61] raise concerns that the sexual rights of WLWH are being unjustly compromised in the current legal climate. While we are unable to directly assess the effect of criminalization of HIV non-disclosure on sexual activity within this study, qualitative studies in Canada have found that fear, anxiety and uncertainty related to the current legal obligation to disclose HIV serostatus to sexual partners have resulted in some PLWH abstaining from sexual activity altogether [70]. Forgoing sexual activity removes expectations and risks of HIV disclosure to sexual partners, difficulties negotiating condom use and risks of sexual HIV transmission. For some WLWH, sexual abstinence may also be a response to fear and threats of rejection by sexual partners, secondary disclosure, relationship dissolution, violence and social isolation following disclosure of HIV status [32,58,71,72].

Despite the high prevalence of sexual inactivity in our cohort, a large proportion of women overall report being

satisfied with their current sex lives (64%). While satisfaction differs by sexual activity, it is higher overall than expected based on previous research [60]. Among sexually inactive women, half report being satisfied with their present sex lives, suggesting that sexual abstinence may be deliberate or intentional, not simply circumstantial. However, sexually inactive women report higher HIV-related stigma. This suggests that HIV-related stigma plays an important role in sexual decision-making, expression and lives of WLWH. Furthermore, WLWH experiencing high HIV-related stigma may be actively choosing sexual inactivity as a strategy to avoid having to navigate disclosure and risk stigma and discrimination in initiating sexual partnerships, and for many women this is a satisfactory approach to their sexual lives.

Interventions are required to reframe the approach to sexual health among WLWH. A component of this reframing is a movement towards models that advocate for Positive Health, Dignity and Prevention frameworks [23,73], which place the person living with HIV at the centre of their health,

care and well-being, well beyond a role in “positive prevention” of on-going transmission of HIV. For example, the Au-delà du VIH: être femme Plurielle [Beyond HIV: valuing the many aspects of women’s lives] program in Quebec is grounded in an empowerment approach to sexual health for WLWH, and supports the capacity of WLWH to respond to their sexual needs and mobilize available resources to gain control over their sexual lives. This, and other similar interventions, recognize WLWH as sexual beings with sexual rights and aim to overcome beliefs and taboos of female sexuality and desire [74].

Limitations to this study are acknowledged. First, cross-sectional analyses preclude determination of causality between correlates and sexual inactivity, and reverse causality is possible (e.g. sexual inactivity leading to depression). Second, there is risk of reporting bias whereby women may underreport sexual activity because of prevailing stigma and discrimination against sexually active WLWH. However, surveys were administered by PRAs, a diverse group of WLWH with identities and life experiences in common with participants, which may have decreased the extent of social desirability bias in reports about sexual activity (with <10% of participants completing the sexual health section independently). Third, we are unable to report whether sexual inactivity represents deliberate or intentional abstinence for sex or a lack of opportunity for sex; however, this difference will be explored in the CHIWOS 18-month follow-up questionnaire. Finally, several important psychosocial aspects of sexual inactivity, including sensation seeking, desire and motivation, as well as knowledge and perceived effect of criminalization of HIV non-disclosure on sexual activity, were not directly assessed in this study, and are likely to be important explanatory pathways to understanding sexual inactivity among WLWH. These pathways will also be assessed in the follow-up questionnaire.

## Conclusions

A satisfying sexual life is a critical component of health and well-being for all people, including WLWH. The observed high rate of sexual inactivity coupled with associations with HIV-related stigma and sexual dissatisfaction underscores a need to revisit the narrative about sexual activity among WLWH. This narrative has focused heavily on HIV risk reduction approaches and now increasingly on the public health importance of viral suppression and the legal test for HIV non-disclosure, but much less has been said about healthy sexuality. A transitioning of the approach to healthy sexuality among WLWH requires that we finally erase the current view of WLWH as “vectors, vessels and victims” of HIV [75], rather as empowered individuals with agency and deserving of intimate relationships. These findings underscore an urgent need for public health and socio-structural interventions to de-stigmatize HIV, support safe disclosure and re-appropriate the sexual rights of WLWH.

### Authors’ affiliations

<sup>1</sup>Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada; <sup>2</sup>British Columbia Centre for Excellence in HIV/AIDS, Vancouver, British Columbia, Canada; <sup>3</sup>Chronic Viral Illness Service, McGill University Health Centre, Montreal, Quebec, Canada; <sup>4</sup>Department of Family Medicine, McGill University, Montreal, Quebec, Canada; <sup>5</sup>Women’s College

Research Institute, Women’s College Hospital, Toronto, Ontario, Canada; <sup>6</sup>Women’s Health in Women’s Hands Community Health Centre, Toronto, Ontario, Canada; <sup>7</sup>Département de Sexologie, Université du Québec à Montréal, Montréal, Quebec, Canada; <sup>8</sup>Department of Medicine and Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada

### Competing interests

The authors have no competing interests to declare.

### Authors’ contributions

AK, AC, AdP, SP, KPB, NO, JTP, KB, VJN, WT, RSH, and ML contributed to the conceptualization and design of the study. AN, PS, and GC conducted the statistical analyses. AK wrote the first draft of the manuscript and all co-authors critically reviewed and revised the manuscript for intellectual content. All authors have read and approved the final version.

### Acknowledgements

The CHIWOS Research Team would like to thank women living with HIV for their contributions to this study. We also thank the national team of co-investigators, collaborators, and Peer Research Associates and acknowledge the national Steering Committee, our three provincial Community Advisory Boards, the National CHIWOS Aboriginal Advisory Board, the BC Centre for Excellence in HIV/AIDS for data support and analysis, and all our partnering organizations for supporting the study.

### Funding sources

CHIWOS is funded by the Canadian Institutes of Health Research (CIHR), the CIHR Canadian HIV Trials Network (CTN 262), the Ontario HIV Treatment Network (OHTN) and the Academic Health Science Centres (AHSC) Alternative Funding Plans (AFP) Innovation Fund. AdP received support from Fonds de Recherche du Québec – Santé (FRQS) (Chercheur-boursier clinicien – Junior 1), SP received support in the form of a Study Abroad Studentship from the Leverhulme Trust, NO received support from FRQS and AK received salary support through a Tier 2 Canada Research Chair in Global HIV and Sexual and Reproductive Health.

**The CHIWOS Research Team:** Rahma Abdul-Noor (*Women’s College Research Institute*), Aranka Anema (*University of British Columbia*), Jonathan Angel (*Ottawa Hospital Research Institute*), Jean-Guy Baril (*Clinique du Quartier-Latin*), Fatimatou Barry (*Women’s College Research Institute*), Greta Bauer (*University of Western Ontario*), Kerrigan Beaver (*Women’s College Research Institute*), Denise Becker (*Positive Living Society of British Columbia*), Anita Benoit (*Women’s College Research Institute*), Jason Brophy (*Children’s Hospital of Eastern Ontario*), Lori Brotto (*University of British Columbia*), Ann Burchell (*Ontario HIV Treatment Network*), Claudette Cardinal (*Simon Fraser University*), Allison Carter (*British Columbia Centre for Excellence in HIV/AIDS and Simon Fraser University*), Angela Cescon (*British Columbia Centre for Excellence in HIV/AIDS*), Lynne Cioppa (*Women’s College Research Institute*), Jeffrey Cohen (*Windsor Regional Hospital*), Guillaume Colley (*British Columbia Centre for Excellence in HIV/AIDS*), Tracey Conway (*Women’s College Research Institute*), Curtis Cooper (*Ottawa Hospital Research Institute*), Jasmine Cotnam (*Women’s College Research Institute*), Janette Cousineau (*Women’s College Research Institute*), Janice Dayle (*McGill University Health Centre*), Marisol Desbiens (*Women’s College Research Institute*), Hania Dubinsky (*McGill University Health Centre*), Danièle Dubuc (*McGill University Health Centre*), Janice Duddy (*Pacific AIDS Network*), Brenda Gagnier (*Women’s College Research Institute*), Jacqueline Gahagan (*Dalhousie University*), Claudine Gasingirwa (*Women’s College Research Institute*), Nada Gataric (*British Columbia Centre for Excellence in HIV/AIDS*), Saara Greene (*McMaster University*), Trevor Hart (*Ryerson University*), Catherine Hankins (*UNAIDS*), Bob Hogg (*British Columbia Centre for Excellence in HIV/AIDS and Simon Fraser University*), Terry Howard (*Positive Living Society of British Columbia*), Shazia Islam (*Women’s College Research Institute*), Evin Jones (*Pacific AIDS Network*), Charu Kaushic (*McMaster University*), Alexandria Keating (*VIVA and Southern Gulf Islands AIDS Society*), Logan Kennedy (*Women’s College Research Institute*), Mary Kestler (*Oak Tree Clinic, BC Women’s Hospital and Health Centre*), Maxime Kiboyogo (*McGill University Health Centre*), Marina Klein (*McGill University Health Centre*), Gladys Kwaramba (*Women’s College Research Institute*), Andrea Langlois (*Pacific AIDS Network*), Rebecca Lee (*CIHR Canadian HIV Trials Network*), Lynne Leonard (*University of Ottawa*), Johanna Lewis (*Women’s College Research Institute*), Viviane Lima (*British Columbia Centre for Excellence in HIV/AIDS*), Elisa Lloyd-Smith (*Providence Health Care*), Carmen Logie (*University of Toronto*), Shari Margolese (*Women’s College Research Institute*), Carrie Martin (*Native Women’s Shelter of Montreal*), Renee



Masching (*Canadian Aboriginal AIDS Network*), Lyne Massie (*Université du Québec à Montréal*), Melissa Medjuck (*Positive Women's Network*), Brigitte Ménard, (*McGill University Health Centre*), Cari Miller (*Simon Fraser University*), Deborah Money (*Women's Health Research Institute*), Marvelous Muchenje (*Women's Health in Women's Hands*), Mary Mwalwanda (*Women's College Research Institute*), Mary (Muthoni) Ndung'u (*Women's College Research Institute*), Valerie Nicholson (*Simon Fraser University*), Illuminée Nzikwizika (*McGill University Health Centre*), Kelly O'Brien (*University of Toronto*), Nadia O'Brien (*McGill University Health Centre and McGill University*), Gina Ogilvie (*British Columbia Centre for Disease Control*), Susanna Ogunnaike-Cooke (*Public Health Agency of Canada*), Joanne Otis (*Université du Québec à Montréal*), Ali Palmer (*Simon Fraser University*), Sophie Patterson (*Simon Fraser University*), Doris Peltier (*Canadian Aboriginal AIDS Network*), Yasmeen (Ashria) Persad (*Women's College Research Institute*), Neora Pick (*Oak Tree Clinic, BC Women's Hospital and Health Centre*), Alie Pierre, (*McGill University Health Centre*), Jeff Powis (*Toronto East General Hospital*), Karène Proulx-Boucher (*McGill University Health Centre*), Corinna Quan (*Windsor Regional Hospital*), Janet Raboud (*Ontario HIV Treatment Network*), Anita Rachlis (*Sunnybrook Health Science Centre*), Edward Ralph (*St. Joseph's Health Care*), Stephanie Rawson, (*Simon Fraser University, BC*), Eric Roth (*University of Victoria*), Danielle Rouleau (*Centre Hospitalier de l'Université de Montréal*), Sean Rourke (*Ontario HIV Treatment Network*), Sergio Rueda (*Ontario HIV Treatment Network*), Mercy Saavedra (*Women's College Research Institute*), Kate Salters (*Simon Fraser University*), Margarine Sanchez (*ViVA and Southern Gulf Islands AIDS Society*), Roger Sandre (*Haven Clinic*), Jacque Sas (*CIHR Canadian HIV Trials Network*), Paul Sereda (*British Columbia Centre for Excellence in HIV/AIDS*), Fiona Smaill (*McMaster University*), Stephanie Smith (*Women's College Research Institute*), Marcie Summers (*Positive Women's Network*), Tsitsi Tigere (*Women's College Research Institute*), Wangari Tharao (*Women's Health in Women's Hands*), Jamie Thomas-Pavanel (*Women's College Research Institute*), Christina Tom (*Simon Fraser University, BC*), Cécile Tremblay (*Centre Hospitalier de l'Université de Montréal*), Benoit Trottier (*Clinique l'Actuel*), Sylvie Trottier (*Centre Hospitalier Universitaire de Québec*), Christos Tsoukas (*McGill University Health Centre*), Sharon Walmsley (*Toronto General Research Institute*), Kath Webster (*Simon Fraser University*), Wendy Wobeser (*Kingston University*), Jessica Yee (*Native Youth Sexual Health Network*), Mark Yudin (*St-Michael's Hospital*), Wendy Zhang (*British Columbia Centre for Excellence in HIV/AIDS*). All other CHIWOS Research Team Members wish to remain anonymous.

#### Disclaimer

The authors alone are responsible for the views expressed in this article and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

#### References

- UNAIDS. The Gap Report 2014. [cited 2015 Sep 14]. Available from: [http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS\\_Gap\\_report\\_en.pdf](http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS_Gap_report_en.pdf)
- Public Health Agency of Canada (2011). Estimates of HIV prevalence and incidence in Canada, 2011. Ottawa: Surveillance and Risk Assessment Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada. [cited 2015 Sep 14]. Available from: <http://www.phac-aspc.gc.ca/aids-sida/publication/surveport/assets/pdf/estimat2011-eng.pdf>
- Samji H, Cescon A, Hogg RS, Modur SP, Althoff KN, Buchacz K, et al. Closing the gap: increases in life expectancy among treated HIV-positive individuals in the United States and Canada. *PLoS One*. 2013;8(12):e81355.
- Hogg RS, Yip B, Chan KJ, Wood E, Craib KJP, O'Shaughnessy MV, et al. Rates of disease progression by baseline CD4 cell count and viral load after initiating triple-drug therapy. *J Am Med Assoc*. 2001;286(20):2568–77.
- Patterson S, Cescon A, Samji H, Chan K, Zhang W, Raboud J, et al. Life expectancy of HIV-positive individuals on combination antiretroviral therapy in Canada. *BMC Infect Dis*. 2015;15:274.
- Forbes JC, Alimenti AM, Singer J, Brophy JC, Bitnun A, Samson LM, et al. A national review of vertical HIV transmission. *AIDS*. 2012;26(6):757–63.
- Castilla J, Del Romero J, Hernando V, Marinovich B, García S, Rodríguez C. Effectiveness of highly active antiretroviral therapy in reducing heterosexual transmission of HIV. *J Acquir Immune Defic Syndr*. 2005;40(1):96–101.
- Weber S, Waldura JF, Cohan D. Safer conception options for HIV serodifferent couples in the United States: the experience of the National Perinatal HIV Hotline and Clinicians' Network. *J Acquir Immune Defic Syndr*. 2013;63(4):e140–1.

- UNAIDS (2014). Post-2015: a global movement for HIV and sexual and reproductive health and rights. Geneva: UNAIDS.
- World Health Organization (2002). Defining sexual health. Report of a Technical Consultation on Sexual Health. Geneva: WHO.
- Salmander Trust (2014). Building a safe house on firm ground: key findings from a global values and preferences survey regarding the sexual and reproductive health and human rights of women living with HIV. Geneva: WHO.
- Gruskin S, Ferguson L. Government regulation of sex and sexuality: in their own words. *Reprod Health Matters*. 2009;17(34):108–18.
- Kennedy CE, Spaulding AB, Brickley DB, Almers L, Mirjahangir J, Packel L, et al. Linking sexual and reproductive health and HIV interventions: a systematic review. *J Int AIDS Soc*. 2010;13(1):26.
- Gay J, Hardee K, Croce-Galis M, Hall C. What works to meet the sexual and reproductive health needs of women living with HIV/AIDS. *J Int AIDS Soc*. 2011;14(1):56.
- Shapiro K, Ray S. Sexual health for people living with HIV. *Reprod Health Matters*. 2007;15(29):67–92.
- Greene S. (M)othering With HIV: Reconstructing & Resisting Health & Social Surveillance. 2nd International Conference for the Social Science and Humanities in HIV; July 9 2013; Paris, France. Session CS51.
- Canadian HIV/AIDS Legal Network. The criminalization of HIV non-disclosure in Canada and internationally. 2014 [cited 2015 Jan 1]. Available from: [http://www.aidslaw.ca/site/wp-content/uploads/2014/09/CriminalInfo2014\\_ENG.pdf](http://www.aidslaw.ca/site/wp-content/uploads/2014/09/CriminalInfo2014_ENG.pdf)
- Montaner JS. Treatment as prevention – a double hat-trick. *Lancet*. 2011;378(9787):208–9.
- The Global Network of People Living with HIV (GNP+) (2010). The Global Criminalization Scan Report 2010. Geneva: GNP+.
- UNAIDS (2014). 90-90-90: an ambitious treatment target to help end the AIDS epidemic. Geneva, Switzerland. [cited 2015 Sep 14]. Available from: [http://www.unaids.org/sites/default/files/media\\_asset/90-90-90\\_en\\_0.pdf](http://www.unaids.org/sites/default/files/media_asset/90-90-90_en_0.pdf)
- Bernard E, Bennett-Carlson R. Criminalization of HIV non-disclosure, exposure and transmission: background and current landscape. Geneva: UNAIDS; 2012 [cited 2015 Sep 14]. Available from: [http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/document/2012/BackgroundCurrentLandscapeCriminalisationHIV\\_Final.pdf](http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/document/2012/BackgroundCurrentLandscapeCriminalisationHIV_Final.pdf)
- Canadian AIDS Society (2011). Positive prevention position statement (March 2011). Vancouver [cited 2015 May 7]. Available from: <http://www.cdnaids.ca/positive-prevention-position-statement-march-2011>
- Global Network of People Living with HIV (GNP+) U (2011). Positive health, dignity and prevention: a policy framework. Vancouver. [cited 2015 May 7]. Available from: [http://www.unaids.org/sites/default/files/media\\_asset/20110701\\_PHDP\\_0.pdf](http://www.unaids.org/sites/default/files/media_asset/20110701_PHDP_0.pdf)
- Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med*. 2011;365(6):493–505.
- Loutfy MR, Wu W, Letchumanan M, Bondy L, Antoniou T, Margolese S, et al. Systematic review of HIV transmission between heterosexual serodiscordant couples where the HIV-positive partner is fully suppressed on antiretroviral therapy. *PLoS One*. 2013;8(2):e55747.
- Rodger A BT, Cambiano V, Vernazza P, Estrada V, Van Lunzen J, Collins S, et al. HIV transmission risk through condomless sex if HIV+ partner on suppressive ART: PARTNER Study. 21st Conference on Retroviruses and Opportunistic Infections (CROI 2014); 2014; Boston, MA. March 3–6, 2014. Abstract 153LB.
- Loureço L, Lima VD, Heath K, Nosyk B, Gilbert M, Colley G, et al. Process monitoring of an HIV treatment as prevention program in British Columbia, Canada. *J Acquir Immune Defic Syndr*. 2014;67(3):e94–109.
- World Health Organization (2015). Maximizing the treatment and prevention potential of antiretroviral drugs: early country experiences towards implementing a treat-all policy. WHO. [cited 2015 Sep 14]. Available from: [http://apps.who.int/iris/bitstream/10665/183599/1/WHO\\_HIV\\_2015.35\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/183599/1/WHO_HIV_2015.35_eng.pdf?ua=1)
- Nosyk B, Montaner JS, Colley G, Lima VD, Chan K, Heath K, et al. The cascade of HIV care in British Columbia, Canada, 1996–2011: a population-based retrospective cohort study. *Lancet Infect Dis*. 2014;14(1):40–9.
- Gardner EM, McLees MP, Steiner JF, del Rio C, Burman WJ. The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clin Infect Dis*. 2011;52(6):793–800.
- Loutfy M, Tyndall M, Baril J-G, Montaner JS, Kaul R, Hankins C. Canadian consensus statement on HIV and its transmission in the context of criminal law. *Can J Infect Dis Med Microbiol*. 2014;25(3):135.

32. World Health Organization (2004). Gender dimensions of HIV status disclosure to sexual partners: rates, barriers and outcomes. A Review Paper. Geneva: WHO.
33. Rouleau G, Côté J, Cara C. Disclosure experience in a convenience sample of Quebec-born women living with HIV: a phenomenological study. *BMC Women's Health*. 2012;12(1):37.
34. Obermeyer CM, Bajjal P, Pegurri E. Facilitating HIV disclosure across diverse settings: a review. *Am J Public Health*. 2011;101(6):1011–23.
35. Adam BD, Corriveau P, Elliott R, Globerman J, English K, Rourke S. HIV disclosure as practice and public policy. *Crit Public Health*. 2015;25(4):386–397.
36. Mahajan AP, Sayles JN, Patel VA, Remien RH, Ortiz D, Szekeres G, et al. Stigma in the HIV/AIDS epidemic: a review of the literature and recommendations for the way forward. *AIDS*. 2008;22(Suppl 2):S67.
37. Lowther K, Selman L, Harding R, Higginson IJ. Experience of persistent psychological symptoms and perceived stigma among people with HIV on antiretroviral therapy (ART): a systematic review. *Int J Nurs Stud*. 2014;51(8):1171–89.
38. BC Centre for Excellence in HIV/AIDS (2013). STOP HIV/AIDS: seek and treat for optimal prevention of HIV/AIDS. Vancouver. [cited 2015 May 7]. Available from: <http://stophivaids.ca>
39. BC Centre for Excellence in HIV/AIDS (2013). It's different now. Vancouver. [cited 2015 May 7]. Available from: <http://www.itsdifferentnowbc.org>
40. Public Health Agency of Canada (2012). HIV and AIDS in Canada: Surveillance Report to December 31, 2012. Ottawa: Surveillance and Risk Assessment Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada.
41. De Reus LA, Few AL, Blume LB. Multicultural and critical race feminisms: theorizing families in the third wave. In: Bengtson VL, Acock AC, Allen KR, Dilworth-Anderson P, Klein DM, editors. *Sourcebook of Family Theory and Research*. Thousand Oaks, CA: SAGE Publications; 2005. p. 447–68.
42. Raphael D. *Social determinants of health: Canadian Perspectives*. 2nd ed. Toronto: Canadian Scholars' Press; 2009.
43. Benoit C, Shumka L. Gendering the population health perspective: fundamental determinants of women's health. Final Report Prepared for the Women's Health Research Network. Women's health research network, Vancouver, BC; 2007.
44. Carter A, Greene S, Nicholson V, O'Brien N, Sanchez M, Pokomandy AD, et al. Breaking the glass ceiling: increasing the meaningful involvement of women living with HIV/AIDS (MIWA) in the design and delivery of HIV/AIDS services. *Health Care Women Int*. 2015;36(8):936–64.
45. Kaida A, Carter A, Lemay J, O'Brien N, Greene S, Nicholson V, et al. Hiring, training, and supporting peer researchers: operationalizing community-based research principles within epidemiological studies by, with, and for women living with HIV. 23rd Canadian Association for HIV Research (CAHR) conference; 2014 May 1–4; St. John's, Newfoundland. (Abstract O106); 2014.
46. Abelsohn K, Benoit AC, Conway T, Cioppa L, Smith S, Kwaramba G, et al. "Hear(ing) new voices": peer reflections from community based survey development with women living with HIV. *Prog Community Health Partnersh*. (In Press).
47. Meston C, Trapnell P. Development and validation of a five-factor sexual satisfaction and distress scale for women: the Sexual Satisfaction Scale for Women (SSS-W). *J Sex Med*. 2005;2(1):66.
48. Carter A, Sereda P, Nohpal A, Colley G, Proulx-Boucher K, Thomas-Pavanel J, et al. Validating self-report of undetectable viral load against laboratory plasma viral load measures in a multi-site cohort of women living with HIV in British Columbia. 24th Canadian Association for HIV Research (CAHR) Conference; April 3–May 3 2015; Toronto, Ontario. (Abstract EPH112); 2015.
49. Revicki DA, Sorensen S, Wu AW. Reliability and validity of physical and mental health summary scores from the Medical Outcomes Study HIV Health Survey. *Med Care*. 1998;36(2):126–37.
50. Wu AW. *MOS-HIV health survey users manual*. Baltimore, MD: Johns Hopkins University; 1999.
51. Zhang W, O'Brien N, Forrest JI, Salters KA, Patterson TL, Montaner JS, et al. Validating a shortened depression scale (10 item CES-D) among HIV-positive people in British Columbia, Canada. *PLoS One*. 2012;7(7):e40793.
52. Radloff LS. The CES-D scale a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1(3):385–401.
53. Wright K, Naar-King S, Lam P, Templin T, Frey M. Stigma scale revised: reliability and validity of a brief measure of stigma for HIV+ youth. *J Adolesc Health*. 2007;40(1):96–8.
54. Berger BE, Ferrans CE, Lashley FR. Measuring stigma in people with HIV: psychometric assessment of the HIV stigma scale. *Res Nurs Health*. 2001; 24(6):518–29.
55. Akaike H. A new look at the statistical model identification. *IEEE Trans Automat Control*. 1974;19(6):716–23.
56. Wilson TE, Jean-Louis G, Schwartz R, Golub ET, Cohen MH, Maki P, et al. HIV infection and women's sexual functioning. *J Acquir Immune Defic Syndr* (1999). 2010;54(4):360–7.
57. Ciccarone DH, Kanouse DE, Collins RL, Miu A, Chen JL, Morton SC, et al. Sex without disclosure of positive HIV serostatus in a US probability sample of persons receiving medical care for HIV infection. *Am J Public Health*. 2003;93(6):949–54.
58. Bogart LM, Collins RL, Kanouse DE, Cunningham W, Beckman R, Golinelli D, et al. Patterns and correlates of deliberate abstinence among men and women with HIV/AIDS. *Am J Public Health*. 2006;96(6):1078.
59. Siegel K, Schrimshaw EW. Reasons for the adoption of celibacy among older men and women living with HIV/AIDS. *J Sex Res*. 2003;40(2):189–200.
60. Shahhosseini Z, Gardeshi ZH, Pourasghar M, Salehi F. A review of affecting factors on sexual satisfaction in women. *Mater Sociomed*. 2014;26(6):378–81.
61. Greene S, Symington A, Ion A, Carvalho A, Loutfy M. Judging mothers: criminalization's creep into the health and social care of HIV-positive mothers. Poster 213. 23rd Canadian Association for HIV Research (CAHR) conference; 2014 May 1–4; St. John's, Newfoundland. (Abstract P213); 2014.
62. Leigh BC, Temple MT, Trocki KF. The sexual behavior of US adults: results from a national survey. *Am J Public Health*. 1993;83(10):1400–8.
63. Siegel K, Schrimshaw EW, Lekas H-M. Diminished sexual activity, interest, and feelings of attractiveness among HIV-infected women in two eras of the AIDS epidemic. *Arch Sex Behav*. 2006;35(4):437–49.
64. Sow K, Desclaux A. Stratégies féminines face au risque de transmission sexuelle du VIH au temps des antirétroviraux. Les femmes à l'épreuve du VIH au Sud. [Women's strategies to counter risk of HIV sexual transmission in the era of antiretrovirals. Women's protection against HIV in the South.] 2011: 165–77.
65. Logie CH, James L, Tharao W, Loutfy MR. HIV, gender, race, sexual orientation, and sex work: a qualitative study of intersectional stigma experienced by HIV-positive women in Ontario, Canada. *PLoS Med*. 2011;8(11): e1001124.
66. Carr RL, Gramling LF. Stigma: a health barrier for women with HIV/AIDS. *J Assoc Nurses AIDS Care*. 2004;15(5):30–9.
67. O'Byrne P, Willmore J, Bryan A, Friedman DS, Hendriks A, Horvath C, et al. Nondisclosure prosecutions and population health outcomes: examining HIV testing, HIV diagnoses, and the attitudes of men who have sex with men following nondisclosure prosecution media releases in Ottawa, Canada. *BMC Public Health*. 2013;13:94.
68. Adam B, Elliott R, Corriveau P, English K. Impacts of criminalization on everyday lives of people living with HIV in Canada. *Sex Res Soc Policy J*. 2014;11:39–49.
69. Mykhalovskiy E, Betteridge G, McLay D. HIV non-disclosure and the criminal law: establishing policy options for Ontario. [Section 4: Social research on the use of the criminal law to address HIV non-disclosure in Ontario: findings from a qualitative research study]. 2010 [cited 2014 Dec 31]. Available from: [http://cocqsida.com/assets/files/2.dossiers/Ontario\\_HIV\\_Non-Disclosure\\_and\\_the\\_Criminal\\_Law.pdf](http://cocqsida.com/assets/files/2.dossiers/Ontario_HIV_Non-Disclosure_and_the_Criminal_Law.pdf)
70. Mykhalovskiy E. The problem of "significant risk": exploring the public health impact of criminalizing HIV non-disclosure. *Soc Sci Med*. 2011;73(5): 668–75.
71. Keegan A, Lambert S, Petrak J. Sex and relationships for HIV-positive women since HAART: a qualitative study. *Aids Patient Care STDS*. 2005;19(10): 645–54.
72. Bott S, Obermeyer CM. The social and gender context of HIV disclosure in sub-Saharan Africa: a review of policies and practices. *SAHARA J*. 2013; 10(Suppl 1):S5–16.
73. UNAIDS. Positive health, dignity and prevention. Operational Guidelines 2013 [cited 2015 Sep 14]. Available from: [http://www.unaids.org/sites/default/files/media\\_asset/20130802\\_Positive\\_Health\\_Dignity\\_Prevention\\_Operational\\_Guidelines\\_0.pdf](http://www.unaids.org/sites/default/files/media_asset/20130802_Positive_Health_Dignity_Prevention_Operational_Guidelines_0.pdf)
74. St-Pierre-Gagné S, Otis J, Massie L, Kéïta DB, Yattassaye A, Dème R, et al. Emotional and sexual well-being of women living with HIV in Quebec: Cultural adaptation and formative evaluation of the program <<PLURIELLES>>. 23rd Canadian Association for HIV Research (CAHR) conference; 2014 May 1–4; St. John's, Newfoundland. (Abstract P250); 2014.
75. Csete J. 'Vectors, vessels and victims': HIV/AIDS and women's human rights in Canada. Toronto: Canadian HIV/AIDS Legal Network; 2005.